

Scotland's Rural College

Species-dependent response to the influence of adaptation length during assay for metabolisable energy of cereal grains employing the difference method

Olukosi, OA; Adedokun, SA; Agboola, JO

Published in:

Animal Feed Science and Technology

DOI:

[10.1016/j.anifeedsci.2017.06.010](https://doi.org/10.1016/j.anifeedsci.2017.06.010)

First published: 24/06/2017

Document Version

Peer reviewed version

[Link to publication](#)

Citation for pulished version (APA):

Olukosi, OA., Adedokun, SA., & Agboola, JO. (2017). Species-dependent response to the influence of adaptation length during assay for metabolisable energy of cereal grains employing the difference method. *Animal Feed Science and Technology*, 231, 111 - 118. <https://doi.org/10.1016/j.anifeedsci.2017.06.010>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Species-dependent response to the influence of adaptation length during assay for
metabolisable energy of cereal grains employing the difference method**

Oluyinka A. Olukosi^{a*}, Sunday A. Adedokun^b and Jeleel O. Agboola^{a1}

^a Monogastric Science Research Centre, Scotland's Rural College, Edinburgh, EH9 3JG, UK

^b Department of Animal and Food Sciences, University of Kentucky, Lexington, KY 40546,

USA

*Corresponding author:

Oluyinka A. Olukosi

Monogastric Science Research Centre, Scotland's Rural College,

Edinburgh, EH9 3JG, U.K.

Email: oluyinka.olukosi@sruc.ac.uk

Tel: +44 1292-525103

¹ Present address: Department of Animal Sciences, Wageningen University, De Elst 1, 6708 WD
Wageningen, The Netherlands.

18 ABSTRACT

19 Three experiments were conducted to determine the influence of varying lengths of adaptation
20 on metabolisable energy (AME and AMEn) content of maize and barley for broilers, turkeys and
21 laying hens using the difference method. Three hundred and twenty-four Cobb 500 male broiler
22 chicks (Experiment 1), 162 BUT 10 male turkey poults (Experiment 2) or 162 Lohmann brown
23 laying hens (Experiment 3) were offered a nutrient-adequate pre-experimental diet for at least 11
24 days. The birds were then allocated to 9 dietary treatments (a 3×3 factorial arrangement of
25 treatments) with each treatment replicated 6 times. The factors were three diet types (based on
26 wheat-soybean meal (WS), maize-wheat-soybean meal (MWS) or barley-wheat soybean meal
27 (BWS)) and three length of adaptation to dietary treatments (10, 7 or 4 d). The WS was the
28 references diet whereas MWS and BWS were the assay diets. The adaptation period
29 corresponded to 10, 7 or 4 days of feeding experimental diets prior to the end of each
30 experiment. Excreta were collected on the last two days of each experiment. The AME of maize
31 and barley in the assay diets, in each experiment, were calculated using difference method. On
32 all the responses considered, there was no significant diet type \times adaptation length interaction in
33 any of the poultry species. Regardless of the poultry species, AME was greater ($P < 0.05$) for
34 maize compared with barley. The AME (MJ/kg) for maize was 13.5, 13.5 and 13.6 for broilers,
35 turkeys and laying hens, respectively whereas the corresponding AME (MJ/kg) of barley was
36 12.2, 11.8 and 12.6, respectively. The effect of adaptation length during AME assay was
37 statistically significant in turkeys, tended to be significant in broilers but not significant for
38 laying hens. It was concluded that irrespective of poultry species, the greater overriding factor
39 affecting AME determined by the difference method is the particular feedstuff being assayed.

However, length of adaptation to experimental diets during the assay becomes more important in birds with relatively physiologically immature digestive tract.

Keywords: adaptation length, broilers, cereal grains, metabolizable energy, turkeys

Abbreviations: AME – apparent metabolisable energy; AMEn – nitrogen-corrected apparent metabolisable energy; BWS – barley wheat soybean meal diet; cEM – coefficient of energy metabolisability; CF – crude fibre; CP – crude protein; DM – dry matter; MWS – maize wheat soybean meal diet; N – nitrogen; WS – wheat soybean meal diet.

1. Introduction

The enactment of various legislations that led to the imposition of ban on some feed ingredients has drastically reduced the available feedstuff for animal nutrition. For instance, the abolishment of feed resources of animal origin due to the incidence of transmissible spongiform encephalopathy in most European countries (Veldkamp and Bosch, 2015). Therefore, the quest for novel feed resources is currently receiving attention more than ever before. To accurately compute the nutritive values of these potential ingredients, it is pertinent to investigate the influence of varying length of adaptation on energy utilisation of feedstuff, in order to determine the optimum adaptation period in studies involving energy utilisation of currently used and novel feed ingredients. This becomes imperative in order to comply with the recent advocacy on precision livestock feeding – which involves feeding of the animals as very close as possible to their requirements. To achieve this, a careful selection of methodology which gives a precise availability of energy in an ingredient should be considered. Poultry diets are formulated based on the AME content and this usually accounts for over 60% of total cost of feeding. So an

accurate determination of available energy is essential for optimising the efficiency of the diet through reduction in feeding cost and excretion of nutrients into the environment.

Over the years, several studies have extensively studied the AME of various feedstuffs (Sibbald, 1975; Sibbald and Price, 1975; Sibbald, 1976; Farrell, 1978; Mollah et al., 1983). However, one prominent method used in these aforementioned studies was the evaluation of available energy from digestibility measurement. There are chiefly two procedures in evaluating digestibility, the total collection method, and the index method (Kong and Adeola, 2014). The total collection method is laborious and greatly relies on the quantitative measurement of feed intake and excreta output, whereas the index method depends solely on accurate chemical analysis of the index compounds. In each of the two methods, the birds are usually adapted to their cages and feed before sample collection. The adaptation usually lasts for about 3-14 days (Kong and Adeola, 2014) during which the animal is adjusted to a feeding level to remove extra work incurred through collection, drying and analysis of orts due to feed refusal by the birds. However, irrespective of the digestibility procedures, the energy contents of a feedstuff could be determined by direct, indirect (difference) or regression methods, depending on the contribution of the basal or test ingredient to the component of interest (Kong and Adeola, 2014).

Additionally, several changes occur in the digestive tracts of the birds which may affect the outcome of the feeding experiments (Zavarize et al., 2012). Feed consumption influenced the development of digestive tract of birds (Gracia et al., 2003), and duodenum develops earlier than the jejunum and the ileum (Uni et al., 1999). This affects the rate of nutrient utilisation during the early stage of life (Uni et al., 1999). However, the intestinal development reached its peak earlier in broilers compared with laying hens (Zavarize et al., 2012). Most of the experiments involving energy utilisation are conducted in two phases (Bolarinwa et al., 2012; Adebisi and Olukosi,

2015; Kong and Adeola, 2016). The birds are usually fed a basal diet for about 14 days (Adeola and Ileleji, 2009; Adebisi and Olukosi, 2015) before assay diets are introduced to them. Thus, there is a feed changeover and consequently a requirement for adaptation to the assay diets prior to samples collection. The optimum length of the adaptation to the assay diet is not clearly defined but it is likely to be affected by the nature of the assay feedstuff. Cereals grains are majorly used as energy sources in poultry diets; however, the difference in their content and type of fibre often influence their utilisation (JøRgensen et al., 1996). The structural variation in dietary fibre sources often affect their nutritional, physical and biological properties (Theander et al., 1989). Therefore, the objective of the current studies is to examine the influence of adaptation period on the AME content of cereal grains having different fibre content and type.

2. Materials and methods

Bird management and sample collection procedures used in the studies were approved, as appropriate, by Scotland's Rural College Animal Experiment Committee or University of Kentucky Institutional Animal Care and Use Committee.

2.1. Animals and management

Three hundred and twenty-four Cobb 500 male broiler chicks (Experiment 1) or 162 BUT 10 male turkey poults (Experiment 2) or 162 Lohmann brown laying hens (Experiment 3) were used for the determination of AME and AMEn contents of maize and barley using the difference method. The broiler chicks and poults were obtained from local hatcheries and fed pre-experimental diets for at least 11 days posthatch. The laying hens were obtained at 32 weeks of age and fed pre-experimental diets for three weeks prior to the start of the experiment. The pre-experimental diets for all experiments were formulated to meet energy and nutrients recommendations in accordance with the specific breed requirements. Birds were reared in

houses with facilities to regulate temperature, light, and humidity. The birds had *ad libitum* access to feed and water throughout the pre- and experimental periods. In each experiment, the birds were allocated to 9 dietary treatments in a randomised complete block design using initial body weight as blocking criterion. The treatments were in a 3 × 3 factorial arrangement. The factors were three diet types based on wheat-soybean meal (WS), maize, wheat-soybean meal (MWS) or barley, wheat-soybean meal (BWS) and three adaptation lengths for adaptation to dietary treatment (10, 7 or 4 days prior to the end of respective experiments). In all the three experiments, each treatment was replicated 6 times. However, experiment 1 contained six birds per replicate, whereas three birds per replicate were used in experiment 2 or 3.

2.2. Diets and sample collection

Three experimental diets were used in each of the three experiments. These diets composed of a wheat-soybean meal referenced diet, containing no maize or barley, and two assay diets containing 300 g/kg maize or barley, respectively. Energy yielding ingredients such as wheat, soybean meal, and soy-oil were substituted with maize or barley in a way their ratios were similar across all the experimental diets. This was necessary to determine the AME and AMEn of maize and barley using the difference method. Titanium dioxide (TiO₂) was added to all diets (5 g/kg diet) as an indigestible marker for determination of AME content by the index method. Experiment 1 diets were offered from d 14 to 24, 17 to 24 or 21 to 24 to give 10-, 7- or 4-d adaptation lengths, respectively. Experiments 2 and 3 diets were offered from d 11 to 21, d 14 to 21 or d 17 to 21 which correspond to 10-, 7- and 4-d adaptation lengths, respectively. The ingredient and chemical composition of the experimental diets used in the three experiments are shown in Table 1. Excreta were collected on the last two days of each experiment, pooled within a cage and dried in a convectional oven.

2.3. Chemical analysis

Samples of diets, wheat, maize, barley, and excreta were ground to pass through 0.5 mm screen using a mill grinder (Retsch ZM 100, F. Kurt Retsch GmbH & Co.KG, Haan, Germany) before chemical analysis. Samples of diets, maize, barley and excreta were analysed for N, GE, DM and TiO₂ where necessary. To determine DM content, samples were dried at 105°C for 24 h using AOAC method 934.01 (AOAC, 2006) in a drying oven (Uniterm, Russel-Lindsey Engineering Ltd., Birmingham, England, UK). The analysis of GE was done using a bomb calorimeter (Parr adiabatic bomb calorimeter, model 6200, parr instruments, Moline, IL, USA) using benzoic acid as a calibration standard. Titanium dioxide was analysed as described by Myers et al. (2004) for Experiment 1 or Short et al. (1996) for Experiments 2 and 3. Nitrogen was determined by the combustion method using AOAC method 968.06 (AOAC, 2006).

2.4. Calculations and Statistical analysis

Coefficient of nitrogen or energy retention (En) was calculated using the index method as described in the equation: $En = 1 - [(Ti/To)X (Eo/Ei)]$; where Ti is the concentration of marker in feed; To is the concentration of marker in the excreta; Eo is the concentration of energy or N in the excreta, and Ei is the concentration of energy or N in the feed.

Metabolizable energy (kcal/g) was calculated using this relation: $AME = GE_i -$

$\left(GE_o \times \left(\frac{T_i}{T_o} \right) \right)$ where GE_i and GE_o are the gross energy (MJ/kg) in feed and excreta,

respectively; Ti and To are the concentration of titanium in the diet and excreta, respectively.

The cEM of test feedstuffs were calculated using the indirect method after correcting for the non-energy yielding fractions in the diets as described by Olukosi and Adeola (2009). The

test ingredient cEM was calculated using the equation: $EM_{ti} = \frac{\{EM_{td} - [EM_{rd} \times (1 - FC_{ti/td})]\}}{FC_{ti/td}}$, where

EM_{ti} is the coefficient of energy metabolizability of the test ingredient, EM_{td} is the coefficient of energy metabolisability of the assay diet, EM_{rd} is the coefficient of energy metabolisability of the reference diet, $FC_{ti/td}$ is the fractional contribution of the test ingredient to the assay diet.

The AME (MJ/kg) of test feedstuffs was calculated using the equation $AME_{ti} = EM_{ti} \times GE_{ti}$, where AME_{ti} and GE_{ti} are the metabolisable and gross energy, respectively, of the test feedstuff. EM_{ti} is as defined previously. To determine the AMEn, the caloric value of 34.39 kJ/g was used (Hill and Anderson, 1958).

Data were analysed using the Generalised Linear Models of IBM SPSS statistical package (version 22). Statistical significance was set at $P \leq 0.05$ for all means comparisons. The digestibility of the diets (both DM and N), dietary AME and AMEn were analysed as a 3×3 factorial of diet type (WS, MWS or BWS) and varying adaptation lengths (10 days, 7 days or 4 days) using ANOVA procedures (presented as supplementary data). Similarly, the AME, AMEn, and cEM of test ingredients were analysed as a 2×3 factorial of cereal grains (maize or barley) and different adaptation lengths (10 days, 7 days or 4 days) using ANOVA procedures. The main effects of diets, test ingredients and adaptation lengths and their possible interactions on AME and AMEn of both diets and cereals, and total tract retention were analysed, and data were presented with both the main and simple effects.

3. Results

The analysed nutrient composition of wheat, barley, and maize used in the current study are shown in Table 2. The CP and ash contents in barley and maize were lower compared with

wheat. Conversely, the crude fat content in maize and barley were higher compared with wheat. The CF content in barley was higher compared to wheat which was higher than that of maize.

The AME, AMEn and cEM of maize and barley in broilers receiving experimental diets for different adaptation lengths are presented in Table 3. There was neither significant cereal grain \times adaptation length interaction nor significant main effect of adaptation length for AME, cEM and AMEn. On the other hand, AME, cEM and AMEn were greater ($P < 0.01$) in maize compared with barley.

The AME, cEM and AMEn of maize and barley in turkeys receiving experimental diets for different adaptation lengths are presented in Table 4. There was no significant cereal grain \times adaptation length interaction on AME, cEM, and AMEn. The main effect of cereal grain was significant ($P < 0.05$) on AME, cEM and AMEn in turkeys being greater ($P < 0.01$) for maize compared with barley. There was significant ($P < 0.05$) main effect of adaptation length on AME and cEM but not on AMEn. The AME was greater ($P < 0.05$) when determined at 4-d compared with at 7-d whereas 10-d was intermediate. cEM was greater ($P < 0.05$) at 10-d and 4-d adaptation lengths compared with at 7-d adaptation length.

The AME, cEM and AMEn of maize and barley in laying hens receiving experimental diets for different adaptation lengths are presented in Table 5. There was no significant cereal grain \times adaptation length interaction ($P > 0.05$) nor significant main effect of adaptation length for AME, cEM and AMEn. Conversely, the main effect of cereal grain was significant for AME, cEM and AMEn. The AME, cEM and AMEn were greater ($P < 0.05$) in maize compared with barley.

4. Discussion

195 The current experiments were carried out to investigate the impact that varying lengths of
196 adaptation length during AME assay, using different poultry species, could have on AME value
197 of two cereal grains with different fibre profiles.

198 The nutrient composition of wheat, barley, and maize used in this current study are
199 comparable with values reported previously (Lin et al., 1987). The results obtained in the current
200 study shows that irrespective of the species of poultry birds, increasing the fibre contents of the
201 diet (via test ingredient) results in lower metabolisability and consequently lower AME content.
202 Furthermore, high-fibre feedstuff led to lower total tract DM and N retention. These could be
203 attributed to limited amounts of endogenous enzymes necessary to enhance release of available
204 energy in the cereals grains.

205 Masey O'Neill et al. (2014) noted that starch supplies the majority of the available energy
206 in cereals grains, however, this is encapsulated in the cell wall which makes it inaccessible to
207 poultry birds due to their inability to produce sufficient amounts of endogenous enzymes in
208 hydrolysing plant materials (Bedford, 1995). Maize contains more starch than barley. Depressed
209 starch digestibility lower metabolisable energy content of cereals (Mollah et al., 1983; Rogel et
210 al., 1987). Additionally, high fibre feedstuffs like barley contain NSP which have been reported
211 to increase the viscosity of the digesta in poultry. The presence of B-glucans in barley primarily
212 increase the viscosity of the digesta (Burnett, 1966). Increased viscosity could reduce the
213 digestibility and availability of nutrients to the animals (Bedford, 1995; Choct et al., 1996;
214 Masey O'Neill et al., 2014). The increased viscosity and bulk of intestinal content impede the
215 rate of diffusion of substrate and consequently hinder their effective interaction at the mucosal
216 surface (Choct et al., 1996).

217 Additionally, the viscosity of digesta could decrease DM intake due to the increase in
218 satiety value. Further calculations in the current study showed that the DM intake of birds on
219 maize-based diets was higher compared with the other two diets. Dietary fibre may lower N
220 retention in poultry birds (Janssen and Carré, 1985; Mateos et al., 2012). As expected, the total N
221 retention of broilers decreased with increasing fibre level in the diet. With the highest total N
222 retention observed in MWS compared with WS and BWS. The NSP content of cereals reduced
223 protein digestibility (Choct and Annison, 1992). On the other hand, increasing the fibre content
224 of the diet through addition of wheat and barley as oppose to maize did not decrease total N
225 retention in laying hens and turkeys. Laying hens retained more N in their body compared to
226 broilers and turkeys.

227 The neutral detergent fibre (NDF) content in barley was nearly three-fold that recorded in
228 maize, which is an indication of the level of fibre present in the two test ingredient. The
229 determined AME (MJ/kg) of barley for broilers (12.2) and laying hens (12.6) in this experiment
230 correspond with the values reported by Choct et al. (2001). However, these values were slightly
231 higher than that observed in turkeys. This variation could be associated with the difference in the
232 physiological age of the birds. Although, turkeys and broilers used in this current experiment
233 were of similar age but broilers have more physiologically matured digestive tract than turkeys.

234 Likewise, adults laying hens were used in this bioassay. Adults birds have the ability to
235 utilise NSP more effectively, thus the higher the NSP level in the grain, the larger the difference
236 in metabolisable energy (Choct et al., 2001). Choct et al. (1996) noted that NSP depress the
237 AME content of feedstuff due to their tendency to inhibit digestion of starch, protein and lipid.
238 The AME content of maize reported for the three poultry species are similar and this could be
239 traced to the less viscous nature of the grain. Choct et al. (2001) argued that the variation in

AME content of cereals in young and mature birds could as well be due to the degree of viscosity of the grain. With the effect more pronounced in high viscous grain (barley) than less viscous ones like maize.

In order to acclimatise the birds to a specific feedstuff or eliminate the possible confounding effects of changing over feeds (from pre-experiment diet to test, or assay, diets) in experiments, it is common practice to adapt the experimental animals to test ingredients or feeds for a certain number of days followed by the actual feeding and samples collection periods. There seems to be no consensus on the actual number of days to be used as adaptation period. During this period, several changes occur in the digestive tract which may eventually influence the results of the study. Adaptation length varies from 3-14 days in most experiments. Steenfeldt et al. (1998) employed 3 days of adaptation when determining the digestibility and AME of wheat-based diet in broilers. Hew et al. (1998) and Adebisi and Olukosi (2015) utilised 7 days of adaptation in determining the AME content of wheat and wheat-DDGS respectively, in broilers. 10-d adaptation period was used by Cowieson and Ravindran (2008) in investigating the effect of exogenous enzymes on energy digestibility of maize-based diets in broilers. Likewise, conventional grower diet was fed for 10-d adaptation period in determining the influence of enzyme supplementation on metabolisable energy of solvent extracted rapeseed and sunflower seed meal in chicken, quail and guinea fowl (Mandal et al., 2005). It is possible however, that adaptation period required may differ depending on fibre type and content of test feedstuff or on the age of the birds or their combination.

The data from the current experiment suggest that adaptation periods influenced the metabolisable energy content of cereal grain and assay diets more markedly in turkeys but to a less extent in broilers and hardly at all in laying hens. However, the effect of adaptation period

on AME content of cereal grain in turkeys was inconsistent with shortening length of adaptation, as the highest determined AME was observed with 10-d and 4-d adaptation period compared with 7-d. In broilers, metabolisable energy content of cereal grain and assay diets only marginally reduced with decreasing length of adaptation period. The determined metabolisable energy of cereal grain and assay diets was similar for the three different adaptation periods in laying hens.

The differences in physiological maturity among the different birds used in the current experiment may partly explain why there were differences in the effect of adaptation length during AME assay in broilers, turkey and laying hens. The data from the current study indicate that the effect of varying length of adaptation was more apparent in birds with less matured digestive system, namely broiler and turkeys, rather than the laying hens. Turkeys are likely less physiologically mature at the same age with broilers in view of their different rates of growth as has been demonstrated between turkeys and duck (Applegate et al., 2005). This may explain why the effect of adaptation length was clearly significant in turkey, tended to be significant for broilers but not significant for laying hens. Uni et al. (1995) demonstrated that villi height and width increased 25 to 100% in all segment of small intestine between 4 and 10 d of age. Additionally, digestive enzymes may be limiting for adequate nutrient utilisation posthatch in chick (Nitsan et al., 1991) and poults (Sell et al., 1991) and these enzymes concentration increase with age. Therefore, during the development and maturation of the digestive system in young poultry, dietary nutrients may be poorly utilised, most especially during the first ten days after hatching (Batal and Parsons, 2002).

The lowest AME values was observed between 4 and 7 d post-hatching in chicks (Zelenka, 1968) and turkeys (Sell, 1996) and this later increased with age. Siregar and Farrell

(1980) found that AME values were not affected by age, whereas Bartov (1988) reported a decrease in AME with birds grew older. Similarly, AMEn value increased from d 0 to d 14 for New Hampshire \times Columbian chicks whereas the increase lasted until d 21 of age with the commercial chick (Batal and Parsons, 2002). Also, there was evidence that AME and AMEn varied with age of birds in the study of Lopez and Leeson (2008). There is also the possibility that stability in the microbial population of the digestive tract with age (Lu et al., 2003) influences how birds at different ages respond to the influence of adaptation length on AME of feedstuffs. The variation in energy content of a single feedstuff determined at different period after hatching corroborate the findings in this experiment where adaptation period influenced the metabolisability of energy in cereal grains and assay diets in turkeys and broilers but not in laying hens.

5. Conclusions

The results of the current study show that irrespective of the species of poultry birds, the greater overriding factor for energy values of feedstuff is the specific ingredient being assayed. However, the influence of adaptation period during AME assay becomes relevant when the assay utilises birds with relatively physiologically immature digestive tract.

References

Adebiyi, A.O., Olukosi. O.A. 2015. Metabolisable energy content of wheat distillers' dried grains with solubles supplemented with or without a mixture of carbohydrases and protease for broilers and turkeys. Poult. Sci. 94, 1270-1276.

306 Adeola, O., Ileleji, K. 2009. Comparison of two diet types in the determination of metabolizable
307 energy content of corn distillers dried grains with solubles for broiler chickens by the
308 regression method. Poult. Sci. 88, 579-585.

309 AOAC. 2006. Official Methods of Analysis. 18th ed. Association of Official Analytical Chemist,
310 Arlington, VA, USA.

311 Applegate, T.J., Karcher, D.M., Lilburn, M.S.. 2005. Comparative development of the small
312 intestine in the turkey poult and Pekin duckling. Poult. Sci. 84, 426-431.

313 Bartov, I. 1988. Effect of age of broiler chicks and method of determination on the metabolizable
314 energy of corn. In: Proc XVIII World's Poultry Congress, Nagoya, Japan. p 787-789.

315 Batal, A.B., Parsons, C. 2002. Effects of age on nutrient digestibility in chicks fed different diets.
316 Poult. Sci. 81, 400-407.

317 Bedford, M. 1995. Mechanism of action and potential environmental benefits from the use of
318 feed enzymes. Anim. Feed Sci. Technol. 53, 145-155.

319 Bolarinwa, O., Olukosi, O.A., Adeola, O. 2012. Metabolizable energy value of porcine meat and
320 bone meal for broiler chickens. Can. J. Anim. Sci. 92, 73-78.

321 Burnett, G. 1966. Studies of viscosity as the probable factor involved in the improvement of
322 certain barleys for chickens by enzyme supplementation. Br. Poult. Sci. 7:55-75.

323 Choct, M., Annison, G. 1992. The inhibition of nutrient digestion by wheat pentosans. Br. J.
324 Nutr. 67, 123-132.

325 Choct, M., Hughes, R., Perez-Maldonado, R., van Barneveld, R. 2001. The metabolisable energy
326 value of sorghum and barley for broilers and layers. Australian Poult. Sci. Symposium
327 Proc. 10, 39-42.

328 Choct, M., Hughes, R.J., Wang, J., Bedford M.R., Morgan, A.J., Annison, G.. 1996. Increased
329 small intestinal fermentation is partly responsible for the anti-nutritive activity of non-
330 starch polysaccharides in chickens. *Br. Poult. Sci* 37, 609-621.

331 Cowieson, A., Ravindran, V. 2008. Effect of exogenous enzymes in maize-based diets varying in
332 nutrient density for young broilers: growth performance and digestibility of energy,
333 minerals and amino acids. *Br. Poult. Sci.* 49, 37-44.

334 Farrell, D. 1978. Rapid determination of metabolisable energy of foods using cockerels. *Br.*
335 *Poult. Sci.* 19, 303-308.

336 Gracia, M., Aranibar, M.J., Lazaro, R., Medel, P., Mateos, G.. 2003. Alpha-amylase
337 supplementation of broiler diets based on corn. *Poult. Sci.* 82, 436-442.

338 Hew, L., Ravindran, V., Mollah, Y., Bryden, W. 1998. Influence of exogenous xylanase
339 supplementation on apparent metabolisable energy and amino acid digestibility in wheat
340 for broiler chickens. *Anim. Feed Sci. Technol.* 75, 83-92.

341 Hill, F.W., Anderson, D.L.. 1958. Comparison of metabolisable energy and productive
342 determinations with growing chicks. *J. Nutr.* 64, 587-603.

343 Janssen, W.M.M.A., Carré, B. 1989. Influence of fibre on digestibility of poultry feeds. In:
344 *Recent Developments in Poultry Nutrition.* 1989. Cole, D.J.A. and W. Haresign (Eds).
345 Butterworths, England. pp. 79 – 93.

346 JøRgensen, H., Zhao, X.Q., Knudsen, K.E.B., Eggum, B.O.. 1996. The influence of dietary fibre
347 source and level on the development of the gastrointestinal tract, digestibility and energy
348 metabolism in broiler chickens. *Br. J. Nutr.* 75, 379-395.

349 Kong, C., Adeola, O. 2014. Evaluation of amino acid and energy utilization in feedstuff for swine
350 and poultry diets. *Asian-Australas. J. Anim. Sci.* 27, 917-925.

351 Kong, C., Adeola, O. 2016. Determination of ileal digestible and apparent metabolizable energy
 352 contents of expeller-extracted and solvent-extracted canola meals for broiler chickens by
 353 the regression method. *SpringerPlus* 5,1.

354 Lin, F., Knabe, D., Tanksley, T. 1987. Apparent digestibility of amino acids, gross energy and
 355 starch in corn, sorghum, wheat, barley, oat groats and wheat middlings for growing pigs.
 356 *J. Anim. Sci.* 64, 1655-1665.

357 Lopez, G., Leeson, S. 2008. Assessment of the nitrogen correction factor in evaluating
 358 metabolizable energy of corn and soybean meal in diets for broilers. *Poult. Sci.* 87, 298-
 359 306.

360 Lu, J., Idris, U., Harmon B., Hofacre, C., Maurer, J.J., Lee, M.D.. 2003. Diversity and succession
 361 of the intestinal bacterial community of the maturing broiler chicken. *Appl. Environ.*
 362 *Microb.* 69, 6816-6824.

363 Mandal, A., Elangovan, A.V., Tyagi, P.K., Johri, A.K., Kaur, S. 2005. Effect of enzyme
 364 supplementation on the metabolisable energy content of solvent-extracted rapeseed and
 365 sunflower seed meals for chicken, guinea fowl and quail. *Br Poul Sci.* 46, 75-79.

366 Masey O'Neill, H., Smith, J., Bedford, M.R. 2014. Multicarbohydrase enzymes for non-
 367 ruminants. *Asian-Australas. J. Anim. Sci.* 27, 290-301.

368 Mateos, G.G., Jiménez-Moreno, E., Serrano, M.P., Lázaro, R.P. 2012. Poultry response to high
 369 levels of dietary fibre sources varying in physical and chemical characteristics. *J. Appl.*
 370 *Poult. Res.* 21, 156-174.

371 Mollah, Y., Bryden, W., Wallis, I., Balnave, D., and Annison, E.. 1983. Studies on low
 372 metabolisable energy wheats for poultry using conventional and rapid assay procedures
 373 and the effects of processing. *Br. Poult. Sci.* 24, 81-89.

374 Myers, W., Ludden, P., Nayigihugu, V., Hess B. 2004. Technical note: A procedure for the
375 preparation and quantitative analysis of samples for titanium dioxide. J. Anim. Sci 82,
376 179-183.

377 Nitsan, Z., Ben Avraham, G., Zoref, Z., Nir, I.. 1991. Growth and development of the digestive
378 organs and some enzymes in broiler chicks after hatching. Br. Poult. Sci. 32, 515-523.

379 Olukosi, O.A., Cowieson, A.J., Adeola, O. 2007. Age-related influence of a cocktail of xylanase,
380 amylase, and protease or phytase individually or in combination in broilers. Poult. Sci.
381 86, 77-86.

382 Olukosi, O. A., Adeola, O. 2009. Estimation of the metabolisable energy content of meat and
383 bone meal for swine. J. Anim. Sci. 87, 2590-2599.

384 Rogel, A., Annison, E., Bryden, W., Balnave, D. 1987. The digestion of wheat starch in broiler
385 chickens. Crop Pasture Sci. 38, 639-649.

386 Sell, J., Angel, C., Piquer, F., Mallarino, E., and Al-Batshan, H.. 1991. Developmental patterns of
387 selected characteristics of the gastrointestinal tract of young turkeys. Poult. Sci. 70, 1200-
388 1205.

389 Sell, J.L. 1996. Physiological limitations and potential for improvement in gastrointestinal tract
390 function of poultry. J. Appl. Poult. Res. 5, 96-101.

391 Short, F.J., Gorton, J.W.P., Boorman, K.N.. 1996. Determination of titanium dioxide added as an
392 inert marker in chicken digestibility studies. J. Anim. Sci. Technol. 59, 215-221.

393 Sibbald, I. 1975. Indirect methods for measuring metabolizable energy in poultry feeds and
394 ingredients. Feedstuffs 47, 22-24.

395 Sibbald, I. 1976. A bioassay for true metabolizable energy in feedingstuffs. Poult. Sci. 55, 303-
396 308.

- Sibbald, I., Price, K.. 1975. Variation in the metabolizable energy values of diets and dietary components fed to adult roosters. Poult. Sci 54, 448-456.
- Siregar, A., Farrell, D.. 1980. A comparison of the energy and nitrogen metabolism of fed ducklings and chickens. Br. Poult. Sci. 21, 213-227.
- Steenfeldt, S., Hammershøj, M., Müllertz, A., Jensen, J.F. 1998. Enzyme supplementation of wheat-based diets for broilers: 2. Effect on apparent metabolisable energy content and nutrient digestibility. Anim. Feed Sci. Technol. 75, 45-64.
- Theander, O., Westerlund, E., Åman, P., Graham, H.. 1989. Plant cell walls and monogastric diets. Anim. Feed Sci. Technol. 23, 205-225.
- Uni, Z., Noy, Y., Sklan, D. 1995. Posthatch changes in morphology and function of the small intestines in heavy-and light-strain chicks. Poult. Sci. 74, 1622-1629.
- Uni, Z., Noy, Y., Sklan, D.. 1999. Posthatch development of small intestinal function in the poult. Poult. Sci. 78, 215-222.
- Veldkamp, T., Bosch, G.. 2015. Insects: a protein-rich feed ingredient in pig and poultry diets. Anim. Frontiers 5, 45-50.
- Zavarize, K., Sartori, J.R., Gonzales, E., Pezzato, A.C. 2012. Morphological changes of the intestinal mucosa of broilers and layers as affected by fasting before sample collection. Rev. Bras. Ciência Avícola 14, 21-25.
- Zelenka, J. 1968. Influence of the age of chicken on the metabolisable energy values of poultry diets. Br. Poult Sci. 9, 135-142.

Acknowledgements

419 Part funding by London Livery – Worshipful Company of Poulterers for this study is
420 acknowledged. The assistance of Derek Brown, Andrew Scott, Irene Yuill, and Jenna Morrison
421 in the care of animals is gratefully acknowledged. Jeleel Opeyemi was a visiting scholar to
422 MSRC in Ayr from July to December 2016 on Erasmus Mundus Scholarship funded by
423 European Commission. Scotland's Rural College is supported by the Scottish government
424 (RESAS).

425 **Table 1**

426 Ingredient and chemical composition (g/kg) of experimental diets to determine apparent metabolisable energy content of maize and barley fed for
427 different adaptation lengths in broilers, turkeys and laying hens.

Item	Broilers			Turkeys			Laying hens		
	1	2	3	1	2	3	1	2	3
Items									
Wheat	602.2	411.9	411.9	492.8	333.4	333.4	654.1	433.4	433.4
Soybean meal	300.0	205.2	205.2	400	270.7	270.7	210	139.1	139.1
Soy oil	47.0	32.2	32.2	35.0	23.7	23.7	25.0	16.6	16.6
Dicalcium phosphate	17.5	17.5	17.5	26	26	26	9.5	9.5	9.5
Limestone	14.0	14.0	14.0	17.7	17.7	17.7	85.0	85.0	85.0
Titanium dioxide ^a	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
Maize	0	300	0	0	300	0	0	300	0
Barley	0	0	300	0	0	300	0	0	300
Vitamin-mineral premix ^b	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
L-Lysine HCl	3.6	3.6	3.6	8.0	8.0	8.0	1.0	1.0	1.0
DL-Methionine	1.9	1.9	1.9	3.9	3.9	3.9	1.4	1.4	1.4
L-threonine	0.7	0.7	0.7	2.6	2.6	2.6	0.8	0.8	0.8
Salt (NaCl)	3.1	3.1	3.1	4.0	4.0	4.0	3.2	3.2	3.2

Total	1000	1000	1000	1000	1000	1000	1000	1000	1000
Analysed energy and nutrient composition ^c									
Dry matter, g/kg	898.9	896.9	903.1	892.2	894.4	894.1	893.0	896.8	896.8
Gross energy, MJ/kg	17.4	17.0	17.0	16.4	15.9	15.8	14.7	14.6	14.6
Crude protein (N x 6.25), g/kg	240.6	195.0	199.4	253.1	195.6	206.3	182.5	137.5	142.5

428

429 ^a Prepared as 1g titanium dioxide added to 4g of ground corn.

430 ^b Vitamin A, 5484 IU; vitamin D3, 2643 ICU; vitamin E, 11 IU; menadione sodium bisulfite, 4.38 mg; riboflavin, 5.49 mg, d-panthothenic acid, 11 mg; niancin,

431 44.1 mg, choline chloride, 771 mg; vitamin B12, 13.2 ug; biotin, 55.2 ug; thiamine mononitrate, 2.2 mg; folic acid, 990 ug; pyridoxine hydrochloride, 3.3 mg;

432 iodine, 1.11 mg; manganese, 66.06 mg; copper, 4.44 mg; iron, 44.1 mg; zinc, 44.1 mg; selenium, 300 ug.

433 ^c Values are means of duplicate analyses.

Table 2

Analysed chemical composition of wheat, barley and maize (g/kg, as-is basis).

Item	Wheat	Barley	Maize
Dry matter	882.3	892.9	871.9
Crude protein (N x 6.25)	159.9	109.3	84.4
Gross Energy (MJ/kg)	-	17.9	18.1
Crude fat	11.7	47.4	31.2
Crude fiber	21.7	38.8	15.2
Ash	19.7	15.6	13.0
Acid detergent fiber	41.1	67.9	23.6
Neutral detergent fiber	273.1	186.5	68.5

451 **Table 3**

452 Apparent metabolisable energy, coefficient of energy metabolisability and nitrogen-correct
 453 metabolisable energy of maize and barley in broilers receiving experimental diets for different
 454 adaptation lengths¹.

Cereal grain	Adaptation length	AME (MJ/kg)	cEM	AMEn (MJ/kg)
Means for main effect of cereal grain				
Maize		13.5	0.747	13.4
Barley		12.2	0.685	12.1
Pooled SEM ²		0.233	0.013	0.240
P-values of diet type		0.001	0.003	0.001
Means for main effect of adaptation length				
	10 days	13.3	0.740	13.2
	7 days	13.0	0.720	12.8
	4 days	12.4	0.688	12.2
Pooled SEM ²		0.285	0.016	0.295
P-values of adaptation length		0.089	0.093	0.062
Means for simple main effects				
Maize	10 days	13.9	0.766	13.8
	7 days	13.5	0.749	13.5
	4 days	13.1	0.727	13.0
Barley	10 days	12.8	0.713	12.7
	7 days	12.4	0.691	12.2

	4 days	11.6	0.650	11.4
Pooled SEM ²		0.403	0.023	0.417
Cereal grain × adaptation length		0.865	0.826	0.885

455 ¹n of 6 for the simple effect, 18 for main effect of adaptation length and 27 for main effect of
 456 cereal grain. AME = Apparent metabolisable energy; cEM = Coefficient of energy
 457 metabolisability; AMEn = Nitrogen-corrected apparent metabolisable energy.

458 ²SEM = Standard error of difference of mean.

459

460 **Table 4**

461 Apparent metabolisable energy, coefficient of energy metabolisability and nitrogen-correct
 462 metabolisable energy of maize and barley in turkeys receiving experimental diets for different
 463 adaptation lengths¹.

Cereal grain	Adaptation length	AME (MJ/kg)	cEM	AMEn (MJ/kg)
Means for main effect of cereal grain				
Maize		13.7	0.756	13.3
Barley		11.6	0.649	11.2
Pooled SEM ²		0.340	0.019	0.369
P-values of diet type		0.001	0.002	0.002
Means for main effect of adaptation length				
	10 days	12.9 ^{ab}	0.716 ^a	12.7
	7 days	11.7 ^b	0.651 ^b	11.4
	4 days	13.3 ^a	0.741 ^a	12.8
Pooled SEM ²		0.446	0.025	0.491
P-values of adaptation length		0.046	0.013	0.073
Means for simple main effects				
Maize	10 days	14.4	0.795	14.2
	7 days	12.7	0.703	12.4
	4 days	13.9	0.771	13.4
Barley	10 days	11.4	0.637	11.2
	7 days	10.7	0.600	10.3

	4 days	12.7	0.710	12.1
Pooled SEM ²		0.552	0.031	0.736
Cereal grain × adaptation length		0.417	0.330	0.371

464 ¹n of 6 for the simple effect, 18 for main effect of adaptation length and 27 for main effect of
 465 cereal grain. AME = Apparent metabolisable energy; cEM = Coefficient of energy
 466 metabolisability; AMEn = Nitrogen-corrected apparent metabolisable energy.

467 ²SEM = Standard error of mean.

468 ^{a-b} Means in the same column but with different superscripts are significantly different (P <
 469 0.05).

470

471 **Table 5**
 472 Apparent metabolisable energy, coefficient of energy metabolisability and nitrogen-correct
 473 metabolisable energy of maize and barley in laying hens receiving experimental diets for
 474 different adaptation lengths¹.

Cereal grain	Adaptation length	AME (MJ/kg)	cEM	AMEn (MJ/kg)
Means for main effect of cereal grain				
Maize		13.6	0.751	13.5
Barley		12.6	0.703	12.5
Pooled SEM ²		0.247	0.014	0.282
P-values of diet type		0.022	0.017	0.029
Means for main effect of adaptation length				
	10 days	13.2	0.734	13.2
	7 days	12.6	0.702	12.5
	4 days	13.4	0.746	13.4
Pooled SEM ²		0.312	0.017	0.354
P-values of adaptation length		0.271	0.164	0.267
Means for simple main effects				
Maize	10 days	13.5	0.745	13.4
	7 days	13.2	0.727	13.0
	4 days	14.1	0.777	14.0
Barley	10 days	12.9	0.722	12.9
	7 days	12.1	0.676	12.0

	4 days	12.8	0.714	12.7
Pooled SEM ²		0.437	0.002	0.503
Cereal grain × adaptation length		0.822	0.740	0.799

475 ¹n of 6 for the simple effect, 18 for main effect of adaptation length and 27 for main effect of
 476 cereal grain. AME = Apparent metabolisable energy; cEM = Coefficient of energy
 477 metabolisability; AMEn = Nitrogen-corrected apparent metabolisable energy.
 478 ²SEM = Standard error of mean.

479

480